

Systemic mast cell activation disease: the role of molecular genetic alterations in pathogenesis, heritability and diagnostics

Britta Haenisch, Markus M.
Nöthen and Gerhard J.
Molderings*

*Institute of Human Genetics, University Hos-
pital of Bonn, Germany*

doi:10.1111/j.1365-2567.2012.03627.x

Received 13 June 2012; revised 13 July 2012;
accepted 17 July 2012.

*Correspondence: Gerhard J. Molderings,
Institute of Human Genetics, University
Hospital of Bonn, Sigmund-Freud-Strasse 25,
D-53127 Bonn, Germany.

Email: molderings@uni-bonn.de

Senior author: Gerhard J. Molderings

Summary

Despite increasing understanding of its pathophysiology, the aetiology of systemic mast cell activation disease (MCAD) remains largely unknown. Research has shown that somatic mutations in kinases are necessary for the establishment of a clonal mast cell population, in particular mutations in the tyrosine kinase Kit and in enzymes and receptors with crucial involvement in the regulation of mast cell activity. However, other, as yet undetermined, abnormalities are necessary for the manifestation of clinical disease. The present article reviews molecular genetic research into the identification of disease-associated genes and their mutational alterations. The authors also present novel data on familial systemic MCAD and review the associated literature. Finally, the importance of understanding the molecular basis of inherited mutations in terms of diagnostics and therapy is emphasized.

Keywords: heritability; mast cell activation disease; mast cell activation syndrome; mastocytosis; molecular genetics.

Introduction

The term systemic mast cell activation disease (MCAD; synonym mast cell disease) refers to disorders characterized by an enhanced release of mast cell mediators sometimes accompanied by accumulation of such dysfunctional mast cells which may or may not be readily histologically detectable (ref. 1 and further references therein). According to the current classification^{1–3} three variants of systemic MCAD are distinguished: systemic mastocytosis (SM), mast cell activation syndrome (MCAS) and mast cell leukaemia (MCL). The first of these, SM, is characterized by specific pathological mutations and immunohistochemical findings, which are known as the World Health Organization criteria.⁴ A diagnosis of MCAS^{1–3,5,6} is assigned to patients who present with multiple mast cell mediator-induced symptoms but who do not fulfil the requirements for a diagnosis of SM, and in whom relevant differential diagnoses have been excluded. The last, MCL, is an aggressive mast cell neoplasm which is defined by increased numbers of mast cells in bone marrow smears ($\geq 20\%$) and the presence of circulating mast cells (reviewed in ref. 4).

Few data are available concerning the prevalence of systemic MCAD. Both SM and MCL are rare disorders.^{4,7} For SM, data from the French mastocytosis network Association Française pour les Initiatives de Recherche sur le Mastocyte et Les Mastocytoses,⁸ the Spanish mastocytosis network Red Española de Mastocitosis,^{9,10} the Italian Mastocytosis Registry,¹¹ and the German Competence Network on Mastocytosis (own unpublished results), suggest a prevalence of at least one in 364 000 in Europe. However, given that these data represent only a proportion of all cases, the true prevalence will be higher. Data from a clinical population suggest that the prevalence of MCL is two orders of magnitude lower than that of SM.¹² In contrast, MCAS seems to be a more common disorder. Evidence has been presented that MCAS may be an underlying cause of various clinical presentations, e.g. in subsets of patients with fibromyalgia¹³ and irritable bowel syndrome.¹⁴ Hence, the prevalence of MCAS is likely to lie within the single-digit percentage range.

The relationship between systemic MCAD and cutaneous mastocytosis (CM, synonyms: paediatric or childhood-onset mastocytosis) remains unclear. Early studies suggested that CM and systemic MCAD were separate dis-

Abbreviations: CM cutaneous mastocytosis; GNNK glycine-asparagine-asparagine-lysine; IL-4 interleukin 4; MCAD mast cell activation disease; MCAS mast cell activation syndrome; MCL mast cell leukaemia; SM systemic mastocytosis

ease entities, because the majority of CM patients were found to lack mutations of the tyrosine kinase *KIT* gene.^{15,16} However, subsequent studies have demonstrated that the frequency of clonal *KIT* mutations is similar in patients with CM, SM and MCAS, and that they are present in up to 86% of patients from each diagnostic group.^{17–20} In addition, several studies have reported the evolution of CM into SM, suggesting that the two disorders are not distinct, but may instead be part of a continuous spectrum of mast cell-related dysfunction.^{18,21–24} However, unless otherwise stated, the present review is confined to discussion of the systemic MCAD variants in the definitions of the current consensus, i.e. SM, MCAS and MCL.

Genetic alterations in patients with systemic MCAD

The establishment of an aberrant clonal mast cell population is dependent upon mutations in genes that encode proteins with a crucial involvement in the regulation of mast cell activity. Those studied best are the mutations occurring in the tyrosine kinase Kit.^{18,25–27} In the majority of patients with systemic MCAD, these mutations are limited to derivatives of haematopoietic stem cells.^{28,29} This indicates that these mutations did not arise during early embryonic development, and were therefore somatic rather than germline. The mutations underlying systemic MCAD drive aberrant mediator production/release with or without readily histologically detectable mast cell accumulation. Mast cell accumulation is due predominantly to a decrease in mast cell apoptosis (refs 30,31 and further references therein). On a limited scale, it is also due to an increase in proliferation.^{32,33} The mutations underlying MCL appear to drive malignant mast cell proliferation in addition to aberrant mediator production and release.^{4,34,35}

A general view on genetic alterations in tyrosine kinase Kit

The type-III transmembrane receptor tyrosine kinase Kit plays a crucial role in the development of mast cells, as well as of haematopoietic progenitor cells, melanocytes, primordial germ cells and the interstitial cells of Cajal (for review, see ref. 36). In humans, the *KIT* gene (previously termed *c-kit*), which encodes the tyrosine kinase Kit, is located in the pericentromeric region of the long arm of chromosome 4 (4q12; http://www.ncbi.nlm.nih.gov/sites/entrez?db=gene&cmd=retrieve&dopt=default&list_uids=3815#ubor2_RefSeq). The human *KIT* gene spans approximately 89 kb and contains 21 exons, which are transcribed/translated into a protein with a molecular mass of 145 000 and a length of 976 amino acids (<http://atlasgeneticsoncology.org/Genes/KITID127.html>). Under physiological conditions, binding of the endogenous Kit ligand, i.e. the stem-cell factor, to the extracellular

domain of the receptor results in receptor dimerization, activation of the intracellular tyrosine kinase domain through autophosphorylation of specific tyrosine residues, and receptor activation (for review, see ref. 37). In normal mature mast cells, activation of Kit signalling through stem-cell factor leads to increased cell proliferation and survival, changes in mast cell migration and adhesion, mast cell degranulation and mediator release.

Various somatic heterozygous Kit alterations (point mutations and deletions/insertions) have been detected in patients with systemic MCAD (Table 1). In any given patient, one Kit alteration or a combination of two or more Kit alterations can occur. For several of these genetic alterations of *KIT* (Table 1, bold typed) it has already been demonstrated that they cause changes in the downstream Kit signalling pathways in mast cells. These convert Kit into a constitutively active, dysregulated, stem-cell-factor-independent tyrosine kinase. Independence of stem-cell factor binding may be explained, at least in part, by kinase activity within unusual mutation-induced subcellular locations, such as the Golgi compartment or the endoplasmic reticulum.^{38,39} Abnormalities in the growth, proliferation and survival of mutated mast cells are mediated by activation of the phosphoinositide 3-kinase/protein kinase B/mammalian Target of Rapamycin pathway. Research suggests that enhancement of mast cell degranulation is mediated by tyrosine phosphorylation of non-T-cell activation linker.⁴⁰ Within this context, it is interesting to note that within any given patient, mutations that are functionally activating in mast cells may be functionally inactivating in other Kit-expressing cells, e.g. in the Cajal cells of the gastrointestinal tract.⁴¹ This may be a result of differences in the signalling cascades. This could explain why a minority of patients with systemic MCAD present with constipation rather than diarrhoea.¹

The described Kit isoforms

Six Kit isoforms exist as a result of the presence of three alternative splicing sites. Each isoform is of differing pathophysiological relevance. An additional glutamine residue at amino acid position 252 is present because of the insertion of an alternative splicing consensus sequence at the 3' end (– 5 nucleotides) of intron 4, at the beginning of the coding sequence of exon 5.²⁵ The two isoforms, i.e. 252Q(+) and 252Q(–), are present in similar numbers in patients with MCAD and healthy volunteers (Figure 1),^{25,26} which suggests that they have no crucial involvement in the induction of an unregulated increase in mast cell activity.

Two further isoforms result from alternative 5' splice donor sites,⁴² and these are characterized by the presence or absence of the tetrapeptide sequence glycine-asparagine-asparagine-lysine (GNNK) in the extracellular part

Table 1. Distribution of alterations in the amino acids sequence of the tyrosine kinase Kit (deduced from the nucleotide sequence of the 21 exons of the gene *KIT*) in mast cells from patients with systemic mast cell activation disease (mast cell activation syndrome and systemic mastocytosis).

Ligand(SCF)- binding domain exons 1–5 aa 1–308	Dimerization domain exons 6–7 aa 309–410	Proteolytic cleavage site exons 8–9 aa 411–513	Membrane- spanning region exon 10 aa 514–549	Auto- inhibition exon 11 aa 550–591	Signal transduction exons 12–21, aa 592–976		
					Kinase domain1 Kinase insert sequence,	Kinase domain 2	C-terminus
W8R ¹	D327N ²	D419H/G ²	<i>Del 510–513</i> ^{1,2,5}	E554K ²	Del 592–626 ¹	F782S ¹	Frequently splicing errors ^{1,2}
C12S ¹	E338K ^{1,2}	419InsFF ³	Del 521 ¹	V559I/A/G ^{6,10}	G658E ²	N787D ¹	
L18P ²	Q346L ¹	Del 419 ³	F522C ⁶	V560G ^{2,6,11}	Y672S ²	H790R ¹	
P31T ²	M351E/I ^{1,2}	Del 417–	V530I ⁷	D572A ³	683 Ins R ²	H802Y ²	
41 Stop codon ¹	F355L ¹	419InsY ³	A533D ^{6,8}		S688L ²	814 Stop codon ²	
E53K ^{1,2}	E359V ¹	C443Y ³	M541L ^{1,2,3,8,9}		S709A ¹	A814V/T ¹³	
Del59–437 ¹	359 Stop codon ¹	S464L ²			<i>Del 715</i> ^{1,2,12}	R815K ⁶	
E73R ¹		475 Stop codon ^{1,2}			E720K ²	Ins V815–I816 ⁶	
T74R ¹		S476I ³			M724I ²	D816V/Y/F/H/	
K116N ²		K484R ²			A736V ¹	I ^{3,6,11}	
V214L ²		ITD501–502 ³			D751Y ¹	I817V ⁶	
<i>252Ins Q</i> ^{1,2}		ITD502–503 ³			D760V ²	D820G ¹⁴	
K259E ¹		ITD505–508 ³			E761K ²	S821F ¹	
H265Q ¹		K509I ^{3,4}			764 Stop codon ²	N822I/K ^{15,16}	
E270K ^{1,2}						A829T ¹	
L276S ¹						830 Stop codon ²	
G286R ²						A837V ¹	
						E839K ⁶	
						L862V ^{1,2}	

¹Ref. 26; ²Ref. 25; ³Ref. 18; ⁴Ref. 68; ⁵Ref. 43; ⁶Ref. 27; ⁷Ref. 74; ⁸Ref. 17; ⁹Ref. 19; ¹⁰Ref. 64; ¹¹Ref. 75; ¹²Ref. 45; ¹³Ref. 76; ¹⁴Ref. 77; ¹⁵Ref. 20; ¹⁶Ref. 78.

Three isoforms resulting from alternative splicing are indicated by italic type. Mutations which have been demonstrated to induce an increased mast cell activity are indicated by bold type. aa, amino acid; SCF, stem cell factor.

of the juxtamembrane region. *In vitro*, these isoforms differ substantially in terms of their functional activities, with the GNNK(–) isoform showing tumorigenic potency.^{43,44} Hence, a predominance of the GNNK(–) isoform may contribute to or reflect a pathological increase in the activation of affected mast cells. Studies comparing MCAD patients with healthy controls have reported significantly stronger expression of the GNNK(–) isoform compared with the GNNK(+) isoform in mast cells from patients with MCAD (Figure 1).²⁶ An expression intensity ratio for GNNK(–)/(+) of $\geq 80\%$ was associated almost exclusively with high symptom intensity. Although the reason for the predominant expression of the GNNK(–) isoform in these patients is unknown, its presence appears to support a diagnosis of MCAD. However, the absence of this predominant expression does not exclude the diagnosis.

The third alternative splicing site in KIT results in the presence or absence of a serine residue at position 715 in the kinase insert region of human Kit. This results from an alternative 3' splice acceptor site usage.⁴⁵ The two iso-

forms S715(–) and S715(+) are co-expressed in similar quantities in patients with MCAD and in healthy controls (Figure 1).^{25,26} This suggests that these isoforms have no substantial involvement in the induction of a pathological increase in mast cell activity.

The mutation Kit^{D816V}

In the mast cells of the majority (> 90%) of patients with SM, mutations in the activation loop of Kit are present (most frequently D816V; reviewed in ref. 27). According to recent studies, the pathological immunohistochemical alterations that constitute the World Health Organization criteria (formation of mast cell clusters; spindle-shaped morphology of mast cells; expression of CD25 in mast cells; marked increase in blood tryptase concentration) seem to be both causally related and specific to the occurrence of a mutation at amino acid position 816 of tyrosine kinase Kit in the affected mast cells.^{4,46,47} During SM progression, the Kit mutant D816V may disappear (ref. 48 and own unpublished observation).




Figure 1. Distribution of the following ratios: (i) of Q252(+)/Q252(–) expression intensity (i.e. the amount of PCR amplification product with and without the additional glutamine residue [Q]); (ii) of the GNNK(–) isoform compared to the GNNK(+) isoform [i.e. the amount of PCR amplification product without and with the tetrapeptide sequence glycine-asparagine-asparagine-lysine (GNNK)]; and (iii) of S715(–)/S715(+) expression intensity (i.e. the amount of PCR amplification product with and without the serine residue [S]). Comparison of the distribution of the ratios between patients with systemic mast cell activation disease (MCAD, filled symbols) and healthy probands (healthy persons, open symbols). Ordinate: Ratio of the expression intensities. The black bars represent the respective mean. $P < 0.05$, Fisher's exact test; n.s., not significant.

The D816V Kit mutation is present to a varying degree in patients with myeloid and lymphoproliferative mastocytosis-associated clonal haematological non-mast cell lineage diseases.^{49,50} The occurrence of this mutation in these diseases points to a common bi-committed neoplastic progenitor clone. In several studies, the *KIT* mutation at codon 816 in mast cells was also detected on CD34⁺ haematopoietic cells, eosinophils, monocytic neutrophil-lineage bone marrow precursor cells, and lymphocytes in around 30% of patients with SM.^{28,51}

Genetic polymorphisms associated with Kit mutations

Kit^{D816V}-positive SM has been associated with a polymorphism in the *IL13* promoter gene (Table 2).⁵² This polymorphism leads to a high transcription rate. Hence, the known proliferation-promoting influence of interleukin-13 (IL-13) on mast cells⁵³ should be enhanced. In accordance with this hypothesis, this polymorphism has been correlated with an elevated serum tryptase level, which is a marker of increased mast cell activation in SM.⁴

Daley *et al.*⁵⁴ reported that the polymorphism Q576R in the cytoplasmic domain of the α -subunit of the IL-4 receptor (Table 2) was found more frequently in SM patients whose disease was limited to skin, and who exhibited lower levels of surrogate disease markers. These data suggest that this polymorphism may mitigate disease expression, which may be a result of an increase in the efficacy of IL-4 at the mutated receptor. In accordance

with this hypothesis, the addition of IL-4 to mast cell cultures has been shown to induce apoptosis.⁵⁵

Finally, associations between SM and polymorphisms in *TET2* (a candidate tumour-suppressor gene), the *TNF α* gene, and the *VEGFA* promoter gene have been reported. No association with SM was found for *IL8* (Table 2).

However, none of the findings from the above mentioned candidate gene studies have been consistently replicated in independent studies, and so these findings cannot be regarded as proven.

Genome-wide gene expression profiling of human mast cells carrying Kit^{D816V}

D'Ambrosio *et al.*⁵⁶ obtained gene expression profiles from bone marrow mononuclear cells of eight patients with SM and the Kit^{D816V} mutation, and compared them with those of five healthy controls. Analysis of global median differential expression defined 130 genes with a known functional product that were differentially and significantly expressed in SM patients compared with controls. This suggests that SM is accompanied by an altered gene expression profile in the bone marrow. The most prominent of these differentially expressed genes was tryptase, which showed a 44–6-fold increase compared with controls. Genes involved in cell proliferation, neoplastic transformation, and apoptosis were also up-regulated. However, these findings require confirmation in experiments on purified mast cells from patients with SM and MCAS.

Table 2. Genes and genetic alterations related to systemic mast cell activation disease.

Gene	Function	Genetic alteration	Cells investigated	Summary of findings	Reference
<i>KIT</i>	Tyrosine kinase Kit	Multiple alterations	Mast cells and multiple other haematopoietic lineages	Clearly associated with pathological mast cell activation	Present paper
<i>JAK2</i>	Janus kinase 2	V617F	BMMC	Associated with pathological mast cell activation	79,80
<i>PDGFRα</i>	Tyrosine kinase platelet-derived growth factor receptor α	Fusion of FIP1L1 gene to <i>PDGFRα</i> gene in exon 12	PBMC; BMMC; multiple other haematopoietic lineages including mast cells	Associated with pathological mast cell activation	81–84
<i>NRAS</i>	Protein with intrinsic GTPase activity	G12D G13D	BMMC	Aggressive systemic mastocytosis	85
<i>RASGRP4</i>	Ras guanyl nucleotide-releasing protein	Failure to remove intron 5	Bone marrow mast cells	Systemic mastocytosis	35
<i>CBL</i>	E3-ligase	Multiple alterations	Retroviral expression of Cbl mutants in transplanted bone marrow in mice	Cytokine-independent mast cell activation	86
<i>HRH4</i>	Histamine H ₄ receptor	Truncated splice variants	Cord blood-derived mast cells; eosinophils; basophils; dendritic cells	Inhibition of full-length histamine-H ₄ -receptor function	87
<i>IL13</i>	Interleukin 13	–1112C/T	PBMC	Association with Kit ^{D816V} -positive systemic mastocytosis	52
<i>IL4</i>	Interleukin 4	Q576R	PBMC	Association with Kit ^{D816V} -positive systemic mastocytosis	54
<i>TET2</i>	Candidate tumour suppressor gene	Multiple	BMMC; cord blood-derived mast cells and eosinophils	Association with Kit ^{D816V} -positive systemic mastocytosis	88,89
<i>TNF</i>	Tumour necrosis factor α	238G/A	No information provided	Association with Kit ^{D816V} -positive systemic mastocytosis	90
<i>VEGFA</i>	Vascular endothelial growth factor	–1154A/G	No information provided	Association with Kit ^{D816V} -positive systemic mastocytosis	91

BMMC, bone marrow mononuclear cells; PBMC, peripheral blood mononuclear cells/leucocytes.

Cytogenetic findings associated with the Kit^{D816V} mutation

In a third of SM patients, genetic instability of the haematopoietic cells is reflected in chromosome abnormalities, as detected during cytogenetic analysis of bone marrow cells.^{57–59} These abnormalities include deletions of chromosomes 5, 7, 11 and 20. No correlation between clinical features and cytogenetic findings has been reported.

Genetic alterations beyond Kit

Research has demonstrated induction of SM and MCAS secondary to mutations in the genes encoding the kinases JAK2 and PDGFR α ; the Src-kinases; and proteins of cellular signal transduction (*RASGRP4*, *CBL*-encoded E3 ligase, histamine H₄ receptor; Table 2, references therein).

Heritability of systemic MCAD

Although numerous reports of familial occurrence and concordant twin pairs have strongly suggested a genetic

basis for CM,^{16,17,60–67} few data are available concerning the genetic contribution to systemic MCAD. To date, systemic MCAD has been assumed to be largely sporadic in nature¹⁵, and only three familial cases of systemic MCAD (two SM and one MCAS) have been reported. In one SM family, the functionally activating point mutation Kit^{K509I} was identified in both affected patients (mother and daughter).⁶⁸ In a second SM family, a functionally activating mutation in Kit exon 8, which resulted in the deletion of codon 419, was detected in the three affected patients (mother, daughter, granddaughter) but not in the healthy sister.⁶⁹ One MCAS family has also been reported in which the father and his two children (daughter and son) presented with varying clinical features.⁷⁰ However, the underlying genetic alterations were not investigated.

Familial clustering of systemic MCAD

Demonstration of familial clustering would be an important step towards defining the genetic contribution to the risk of systemic MCAD. However, at the time of writing, no systematic investigation of the heritability of systemic

MCAD has been reported. Therefore, the present authors analysed data from 120 Caucasian index patients (Figure 2) with MCAS ($n = 102$) or SM ($n = 18$), who presented to our research group between May 2005 and November 2010, to determine familial aggregation. A sample of 258 probands was used as a control group representative of the German population. These individuals had been randomly recruited from the German inhabitants of the city of Bonn. Analysis of our data revealed that irrespective of systemic MCAD variant and gender, around 75% of our index patients had at least one first-degree relative with systemic MCAD (Figure 2). Within any given family, the disease variant, the severity of the mediator-related symptoms, and the age at clinical onset differed between individuals. Similar findings were reported in abstracts by Burks *et al.* 2005 and Bursztein *et al.* 2009.^{70,71} The prevalence of systemic MCAD among the first-degree relatives in our sample was 33% (Figure 2), which differed significantly ($P < 0.0001$; Fisher's exact test) from the prevalence in the control group (around 14%; Figure 2). The age distribution was similar in both groups. In both patients and controls, a female preponderance was observed (Figure 2). Our data suggest that systemic MCAD pedigrees include more systemic MCAD cases than would be expected by chance. This finding is consistent with the hypothesis that systemic MCAD has a heritable component, at least in a considerable number of patients.

At the time of writing, the molecular processes that result in a familial aggregation of MCAD remain speculative. The detection of differing systemic MCAD-associated somatic mutations within given families (refs 18,72; own unpublished data) is compatible with the hypothesis that disease arises secondary to a dysfunction of mutated, and as yet unidentified, operator or regulator genes. This dysfunction could result in mutations in *KIT* and other mast-cell-regulatory genes. Within this context, it is interesting to note that most of the insertions and deletions in *KIT* in patients with MCAS involve intron–exon junctions, suggesting that distortion of alternative splicing may contribute to the generation of these novel transcripts.²⁶

Conclusions and perspective

Despite increasing understanding of its pathophysiology, the aetiology of MCAD remains largely unknown. The establishment of an aberrant clonal mast cell population is dependent upon mutations in kinases (particularly in the tyrosine kinase Kit) and in enzymes and receptors crucially involved in the regulation of mast cell activity. Other genetic alterations, some of which are as yet undetermined, appear to be related to systemic MCAD phenotype and prognosis.

Recent findings suggest that the various systemic MCAD variants and clinical phenotypes do not represent distinct disease entities, but rather varying presentations of a




Figure 2. Characteristics of the presently investigated population in terms of the familial aggregation of systemic mast cell activation disease (MCAD). In the index patients, the diagnosis was assigned according to the criteria for MCAD.¹ Due to the anonymous nature of the survey of the control population, clinical diagnoses were assigned in the control group on the basis of a self-report checklist and the exclusion of relevant differential diagnoses (for the checklist, see ref. 92). Asterisks: number of affected persons/total number of persons in the respective group (percentage). A sample of 258 probands randomly recruited from the German inhabitants of the city of Bonn was used as a control group representative of the German population.

common generic root process of mast cell dysfunction.^{2,7,25,26} Interestingly, in contrast to adult-onset systemic MCAD, more than 50% of paediatric cases of cutaneous mastocytosis appear to enter long-term remission spontaneously,^{23,65} though whether such remissions are permanent or relapse in adulthood as systemic MCAD is unknown. The reason for this difference in prognosis – if both are clonal diseases with common roots – awaits clarification. Improved understanding of the biochemical processes leading to this resolution may open new avenues in the treatment of adult systemic MCAD.

To optimize research into the genetic basis of MCAD, a number of approaches may be suitable. First, next-generation sequencing could facilitate identification of the putative mutated disease-related operator and regulator genes in familial cases. Second, studies of animal models displaying disturbances secondary to primary mast cell dysfunction will implicate further genes of relevance to systemic MCAD. In dogs, mast cell tumours are the second most common form of neoplastic disease (7–21% of all cutaneous neoplasms). In up to 40% of all canine mast cell tumours, *KIT* mutations are present. Well-documented breed predispositions indicate an underlying heritable component (for review, see ref. 73). Third, genome-wide association studies could allow the unbiased and systematic identification of systemic MCAD risk genes. Since their introduction, genome-wide association studies have identified risk genes for several multifactorial diseases (<http://www.genome.gov/gwastudies/>). However, these studies showed that most risk genes have only moderate effect sizes, and that analysis of large samples is necessary. Given the high prevalence of the MCAS variant, the recruitment of such samples should be possible at least for MCAS.

Taken together, improved knowledge of the genetic aetiology of MCAD may facilitate the development of new therapeutic strategies, which possibly may also imply curative in addition to symptomatic treatment options.

Acknowledgements

We thank Dr Christine Schmäler for her revision of the manuscript. This study was supported by grants from the B. Braun-Stiftung, Germany and the Förderclub Mastzellforschung e.V., Germany.

Disclosures

The authors declare that they have no competing interests.

Authors' contribution

All authors contributed equally to conception, design and drafting of the manuscript. All authors read and approved the final manuscript.

References

- Molderings GJ, Brettner S, Homann J, Afrin LB. Mast cell activation disease: a concise practical guide for diagnostic workup and therapeutic options. *J Hematol Oncol* 2011; **4**:10.
- Akin C, Valent P, Metcalfe DD. Mast cell activation syndrome: proposed diagnostic criteria. *J Allergy Clin Immunol* 2010; **126**:1099–104.
- Valent P, Akin C, Arock M *et al.* Definitions, Criteria and Global Classification of Mast Cell Disorders with Special Reference to Mast Cell Activation Syndromes: a Consensus Proposal. *Int Arch Allergy Immunol* 2012; **157**:215–25.
- Valent P, Akin C, Escribano L *et al.* Standards and standardization in mastocytosis: consensus statements on diagnostics, treatment recommendations and response criteria. *Eur J Clin Invest* 2007; **37**:435–53.
- Alvarez-Twose I, González de Olano D, Sánchez-Muñoz L *et al.* Clinical, biological, and molecular characteristics of clonal mast cell disorders presenting with systemic mast cell activation symptoms. *J Allergy Clin Immunol* 2010; **125**:1269–78.
- Hamilton MJ, Hornick JL, Akin C, Castells MC, Greenberger NJ. Mast cell activation syndrome: a newly recognized disorder with systemic clinical manifestations. *J Allergy Clin Immunol* 2011; **128**:147–52.
- Hermine O, Lortholary O, Leventhal PS *et al.* Case-control cohort study of patients' perceptions of disability in mastocytosis. *PLoS One* 2008; **3**:e2266.
- Lanternier F, Cohen-Akenine A, Palmerini F *et al.* Phenotypic and genotypic characteristics of mastocytosis according to the age of onset. *PLoS One* 2008; **3**:e1906.
- Escribano L, Alvarez-Twose I, Sánchez-Muñoz L *et al.* Prognosis in adult indolent systemic mastocytosis: a long-term study of the Spanish Network on Mastocytosis in a series of 145 patients. *J Allergy Clin Immunol* 2009; **124**:514–21.
- Alvarez-Twose I, González-de-Olano D, Sánchez-Muñoz L *et al.* Validation of the REMA Score for Predicting Mast Cell Clonality and Systemic Mastocytosis in Patients with Systemic Mast Cell Activation Symptoms. *Int Arch Allergy Immunol* 2011; **157**:275–80.
- Merante S, Magliacane D, Neri I *et al.* The New Italian Mastocytosis Registry. 53rd ASH Annual Meeting and Exposition 2010; Poster 3805, [<http://ash.confex.com/ash/2010/webprogram/Paper28567.html>]
- Lim KH, Tefferi A, Lasho TL *et al.* Systemic mastocytosis in 342 consecutive adults: survival studies and prognostic factors. *Blood* 2009; **113**:5727–36.
- Lucas HJ, Brauch CM, Settla L, Theoharides TC. Fibromyalgia – new concepts of pathogenesis and treatment. *Int J Immunopathol Pharmacol* 2006; **19**:5–9.
- Frieling T, Meis K, Kolck UW *et al.* Evidence for mast cell activation in patients with therapy-resistant irritable bowel syndrome. *Z Gastroenterol* 2011; **49**:191–4.
- Longley BJ, Metcalfe DD, Sharp M, Wang X, Tyrrell L, Lu SZ, Heitjan D, Ma Y. Activating and dominant inactivating c-kit catalytic domain mutations in distinct clinical forms of human mastocytosis. *Proc Natl Acad Sci U S A* 1999; **96**:1609–14.
- Rosbotham JL, Malik NM, Syrris P *et al.* Lack of c-kit mutation in familial urticaria pigmentosa. *Br J Dermatol* 1999; **140**:849–52.
- Tang X, Boxer M, Drummond A, Ogston P, Hodgins M, Burden AD. A germline mutation in KIT in familial diffuse cutaneous mastocytosis. *J Med Genet* 2004; **41**:e88.
- Bodemer C, Hermine O, Palmérini F *et al.* Pediatric mastocytosis is a clonal disease associated with D(816)V and other activating c-KIT mutations. *J Invest Dermatol* 2010; **130**:804–15.
- Foster R, Byrnes E, Meldrum C *et al.* Association of paediatric mastocytosis with a polymorphism resulting in an amino acid substitution (M541L) in the transmembrane domain of c-KIT. *Br J Dermatol* 2008; **159**:1160–9.
- Wasag B, Niedoszytko M, Piskorz A *et al.* Novel activating KIT-N822I mutation in familial cutaneous mastocytosis. *Exp Hematol* 2011; **39**:859–65.
- Noack F, Escribano L, Sotlar K, Nunez R, Schuetze K, Valent P, Horny HP. Evolution of urticaria pigmentosa into indolent systemic mastocytosis: abnormal immunophenotype of mast cells without evidence of c-KIT mutation ASP-816-VAL. *Leuk Lymphoma* 2003; **44**:313–9.
- Vogel NM, Lichtin AE, Hsieh FH. Longitudinal disease progression in mastocytosis syndromes: a retrospective chart review. *J Allergy Clin Immunol* 2006; **117**(Suppl): S125.
- Uzzaman A, Maric I, Noel P, Kettelhut BV, Metcalfe DD, Carter MC. Pediatric-onset mastocytosis: a long term clinical follow-up and correlation with bone marrow histopathology. *Pediatr Blood Cancer* 2009; **53**:629–34.
- Bhamidimarri K, Sidhu L, Xie J, Pulakhandam U, Gumstae V. Gastrointestinal manifestations in cutaneous mastocytosis evolving into systemic mastocytosis. *Am J Gastroenterol* 2010; **105**(Suppl. 1):S348.
- Molderings GJ, Kolck UW, Scheurlen C, Brüss M, Homann J, von Kügelgen I. Multiple novel alterations in Kit tyrosine kinase in patients with gastrointestinally pronounced systemic mast cell activation disorder. *Scand J Gastroenterol* 2007; **42**:1045–53.
- Molderings GJ, Meis K, Kolck UW, Homann J, Frieling T. Comparative analysis of mutation of tyrosine kinase kit in mast cells from patients with systemic mast cell activation syndrome and healthy subjects. *Immunogenetics* 2010; **62**:721–7.

- 27 Orfao A, Garcia-Montero AC, Sanchez L, Escribano L; REMA. Recent advances in the understanding of mastocytosis: the role of KIT mutations. *Br J Haematol* 2007; **138**: 12–30.
- 28 Garcia-Montero AC, Jara-Acevedo M, Teodosio C et al. KIT mutation in mast cells and other bone marrow hematopoietic cell lineages in systemic mast cell disorders: a prospective study of the Spanish Network on Mastocytosis (REMA) in a series of 113 patients. *Blood* 2006; **108**:2366–72.
- 29 Kristensen T, Broesby-Olsen S, Vestergaard H, Bindslev-Jensen C, Møller MB; on behalf of the Mastocytosis Centre Odense University Hospital (MastOUH). Circulating KIT D816V mutation-positive non-mast cells in peripheral blood are characteristic of indolent systemic mastocytosis. *Eur J Haematol* 2012; **89**:42–6.
- 30 Kohno M, Yamasaki S, Tybulewicz VLJ, Saito T. Rapid and large amount of autocrine IL-3 production is responsible for mast cell survival by IgE in the absence of antigen. *Blood* 2005; **105**:2059–65.
- 31 Aichberger KJ, Gleixner KV, Mirkina I et al. Identification of proapoptotic Bim as a tumor suppressor in neoplastic mast cells: role of KIT D816V and effects of various targeted drugs. *Blood* 2009; **114**:5342–51.
- 32 Rottem M, Okada T, Goff JP, Metcalfe DD. Mast cells cultured from the peripheral blood of normal donors and patients with mastocytosis originate from a CD34⁺/FcεRI-cell population. *Blood* 1994; **84**:2489–96.
- 33 Jordan JH, Jäger E, Sperr WR et al. Numbers of colony-forming progenitors in patients with systemic mastocytosis: potential diagnostic implications and comparison with myeloproliferative disorders. *Eur J Clin Invest* 2003; **33**:611–8.
- 34 Ning ZQ, Li J, Arceci RJ. Signal transducer and activator of transcription 3 activation is required for Asp816 mutant c-kit-mediated cytokine-independent survival and proliferation in human leukemia cells. *Blood* 2001; **97**:3559–67.
- 35 Yang Y, Li L, Wong GW, Krilis SA, Madhusudhan MS, Sali A, Stevens RL. RasGRP4, a new mast cell-restricted Ras guanine nucleotide-releasing protein with calcium- and diacylglycerol-binding motifs. Identification of defective variants of this signaling protein in asthma, mastocytosis, and mast cell leukemia patients and demonstration of the importance of RasGRP4 in mast cell development and function. *J Biol Chem* 2002; **277**:25756–74.
- 36 Roskoski R. Structure and regulation of Kit protein-tyrosine kinase – the stem cell factor receptor. *Biochem Biophys Res Commun* 2005; **338**:1307–15.
- 37 Roskoski R. Signaling by Kit protein-tyrosine kinase – the stem cell factor receptor. *Biochem Biophys Res Commun* 2005; **337**:1–13.
- 38 Bougherara H, Subra F, Crépin R, Tauc P, Auclair C, Poul MA. The aberrant localization of oncogenic kit tyrosine kinase receptor mutants is reversed on specific inhibitory treatment. *Mol Cancer Res* 2009; **7**:1525–33.
- 39 Toffalini F, Demoulin JB. New insights into the mechanisms of hematopoietic cell transformation by activated receptor tyrosine kinases. *Blood* 2010; **116**:2429–37.
- 40 Gilfillan AM, Rivera J. The tyrosine kinase network regulating mast cell activation. *Immunol Rev* 2009; **228**:149–69.
- 41 Breuer C, Oh J, Molderings GJ, Schemann M, Kuch B, Mayatepek E, Adam R. Therapy-refractory gastrointestinal motility disorder in a child with c-kit mutations. *World J Gastroenterol* 2010; **16**:4363–6.
- 42 Hayashi S, Kunisada T, Ogawa M, Yamaguchi K, Nishikawa S. Exon skipping by mutation of an authentic splice site of c-kit gene in W/W mouse. *Nucleic Acids Res* 1991; **19**:1267–71.
- 43 Caruana G, Cambareri AC, Ashman LK. Isoforms of c-KIT differ in activation of signalling pathways and transformation of NIH3T3 fibroblasts. *Oncogen* 1999; **18**:5573–81.
- 44 Voytyuk O, Lennartsson J, Mogi A, Caruana G, Courtneidge S, Ashman LK, Rönnstrand L. Src family kinases are involved in the differential signaling from two splice forms of c-Kit. *J Biol Chem* 2003; **278**:9159–66.
- 45 Crosier PS, Ricciardi ST, Hall LR, Vitas MR, Clark SC, Crosier KE. Expression of isoforms of the human receptor tyrosine kinase c-kit in leukemic cell lines and acute myeloid leukemia. *Blood* 1993; **82**:1151–8.
- 46 Mayerhofer M, Gleixner KV, Hoelbl A et al. Unique effects of KIT D816V in BaF3 cells: induction of cluster formation, histamine synthesis, and early mast cell differentiation antigens. *J Immunol* 2008; **180**:5466–76.
- 47 Yang Y, Létard S, Borge L. Pediatric mastocytosis-associated KIT extracellular domain mutations exhibit different functional and signaling properties compared with KIT-phosphotransferase domain mutations. *Blood* 2010; **116**:1114–23.
- 48 Gotlib J, Berubé C, Growney JD et al. Activity of the tyrosine kinase inhibitor PKC412 in a patient with mast cell leukemia with the D816V KIT mutation. *Blood* 2005; **106**:2865–70.
- 49 Fritsche-Polanz R, Fritz M, Huber A, Sotlar K, Sperr WR, Mannhalter C, Födinger M, Valent P. High frequency of concomitant mastocytosis in patients with acute myeloid leukemia exhibiting the transforming KIT mutation D816V. *Mol Oncol* 2010; **4**:335–46.
- 50 Sotlar K, Colak S, Bache A, Berezowska S, Krokowski M, Bültmann B, Valent P, Horny HP. Variable presence of KIT(D816V) in clonal haematological non-mast cell lineage diseases associated with systemic mastocytosis (SM-AHNMD). *J Pathol* 2010; **220**: 586–95.
- 51 Akin C, Kirshenbaum AS, Semere T, Worobec AS, Scott LM, Metcalfe DD. Analysis of the surface expression of c-kit Asp816Val activating mutation in T cells, B cells, and myelomonocytic cells in patients with mastocytosis. *Exp Hematol* 2000; **28**:140–7.
- 52 Nedoszytko B, Nedoszytko M, Lange M et al. Interleukin-13 promoter gene polymorphism –1112C/T is associated with the systemic form of mastocytosis. *Allergy* 2009; **64**:287–94.
- 53 Kaur D, Hollins F, Woodman L, Yang W, Monk P, May R, Bradding P, Brightling CE. Mast cells express IL-13Rα1: IL-13 promotes human lung mast cell proliferation and FcεRI expression. *Allergy* 2006; **61**:1047–53.
- 54 Daley T, Metcalfe DD, Akin C. Association of the Q576R polymorphism in the interleukin-4 receptor α chain with indolent mastocytosis limited to the skin. *Blood* 2001; **98**:880–2.
- 55 Bailey DP, Kashyap M, Mirmonsef P, Bouton LA, Domen J, Zhu J, Dessypris EN, Ryan JJ. Interleukin-4 elicits apoptosis of developing mast cells via a Stat6-dependent mitochondrial pathway. *Exp Hematol* 2004; **32**:52–9.
- 56 D'Ambrosio C, Akin C, Wu Y, Magnusson MK, Metcalfe DD. Gene expression analysis in mastocytosis reveals a highly consistent profile with candidate molecular markers. *J Allergy Clin Immunol* 2003; **112**:1162–70.
- 57 Swolin B, Rödger S, Roupe G. Cytogenetic Studies in Patients with Mastocytosis. *Cancer Genet Cytogenet* 2000; **120**:131–5.
- 58 Swolin B, Rödger S, Ögård I, Roupe G. Trisomies 8 and 9 not detected with fish in patients with mastocytosis. *Am J Hematol* 2002; **70**:324–5.
- 59 Jost E, Michaux L, van den Abeele M et al. Complex karyotype and absence of mutation in the c-Kit receptor in aggressive mastocytosis presenting with pelvic osteolysis, eosinophilia and brain damage. *Ann Hematol* 2001; **80**:302–7.
- 60 Offidani A, Cellini A, Simonetti O, Bossi G. Urticaria pigmentosa in monozygotic twins. *Arch Dermatol* 1994; **130**:935–6.
- 61 Pec J, Palencarova E, Malisova S, Dobrota D, Hajtman A, Pec M, Lepej J. Urticaria pigmentosa in identical male twins. *Acta Derm Venereol* 1995; **75**:244.
- 62 Nishida T, Hirota S, Taniguchi M et al. Familial gastrointestinal stromal tumours with germline mutation of the KIT gene. *Nat Genet* 1998; **19**:323–4.
- 63 Trevisan G, Pauluzzi P, Gatti A, Semeraro A. Familial mastocytosis associated with neurosensory deafness. *J Eur Acad Dermatol Venereol* 2000; **14**:119–22.
- 64 Beghini A, Tibiletti MG, Roversi G, Chiaravalli AM, Serio G, Capella C, Larizza L. Germline mutation in the juxtamembrane domain of the kit gene in a family with gastrointestinal stromal tumors and urticaria pigmentosa. *Cancer* 2001; **92**:657–62.
- 65 Ben-Amitai D, Metzker A, Cohen HA. Pediatric cutaneous mastocytosis: a review of 180 patients. *Isr Med Assoc J* 2005; **7**:320–2.
- 66 Yanagihori H, Oyama N, Nakamura K, Kaneko F. C-kit mutations in patients with childhood-onset mastocytosis and genotype-phenotype correlation. *J Mol Diagn* 2005; **7**:252–7.
- 67 de la Sotta PD, Romero WA, Kramer D, Cárdenas C, González S. Cutaneous mastocytosis in twins: multiple mastocytomas and urticaria pigmentosa in two pairs of monozygotic twins. *Pediatr Dermatol* 2011; **28**:585–7.
- 68 Zhang LY, Smith ML, Schultheis B, Fitzgibbon J, Lister TA, Melo JV, Cross NCP, Cavenagh JD. A novel K5091 mutation of KIT identified in familial mastocytosis – *in vitro* and *in vivo* responsiveness to imatinib therapy. *Leuk Res* 2006; **30**:373–8.
- 69 Hartmann K, Wardelmann E, Ma Y et al. Novel germline mutation of KIT associated with familial gastrointestinal stromal tumors and mastocytosis. *Gastroenterology* 2005; **129**:1042–6.
- 70 Burks KD, Wenzel SE, Jones SM. Familial cases of mast cell diseases[abstract]. *J Invest Med* 2005; **53**:S300.
- 71 Bursztejn AC, Bronner M, Kanny G, Barbaud A, Morisset M, Jonveaux P. Genetic evaluation of familial mastocytosis[abstract]. *J Invest Dermatol* 2009; **129**:2528.
- 72 Verzijl A, Heide R, Oranje AP, van Schaik RHN. c-Kit Asp-816-Val mutation analysis in patients with mastocytosis. *Dermatology* 2007; **214**:15–20.
- 73 Welle MM, Bley CR, Howard J, Rüfenacht S. Canine mast cell tumours: a review of the pathogenesis, clinical features, pathology and treatment. *Vet Dermatol* 2008; **19**:321–39.
- 74 O'Brien S, Tefferi A, Valent P. Chronic myelogenous leukemia and myeloproliferative disease. *Hematology Am Soc Hematol Educ Program* 2004; 146–62.
- 75 Buttner C, Henz BM, Welker P, Sepp NT, Grabbe J. Identification of activating c-kit mutations in adult-, but not in childhood-onset indolent mastocytosis: a possible explanation for divergent clinical behavior. *J Invest Dermatol* 1998; **111**:1227–31.
- 76 Sotlar K, Horny HP, Simonitsch I et al. CD25 indicates the neoplastic phenotype of mast cells: a novel immunohistochemical marker for the diagnosis of systemic mastocytosis (SM) in routinely processed bone marrow biopsy specimens. *Am J Surg Pathol* 2004; **28**:1319–25.
- 77 Pignon JM, Giraudier S, Duquesnoy P, Jouault H, Imbert M, Vainchenker W, Vernant JP, Tulliez M. A new c-kit mutation in a case of aggressive mast cell disease. *Br J Haematol* 1997; **96**:374–6.
- 78 Baek JO, Kang HK, Na SY, Lee JR, Roh JY, Lee JH, Kim HJ, Park S. N822K c-kit mutation in CD30-positive cutaneous pleomorphic mastocytosis after germ cell tumour of the ovary. *Br J Dermatol* 2012; **166**:1370–3.

- 79 Galderisi CD, Corless CL, Wolford J, Harrell T, Heinrich MC, Press RD. Simultaneous JAK2 V617F and KIT D816V mutations in systemic mastocytosis. *Mod Pathol* 2006; **19**:225A–6A.
- 80 Steensma DP, Dewald GW, Lasho TL, Powell HL, McClure RF, Levine RL, Gilliland DG, Tefferi A. The JAK2 V617F activating tyrosine kinase mutation is an infrequent event in both "atypical" myeloproliferative disorders and myelodysplastic syndromes. *Blood* 2005; **106**:1207–9.
- 81 Cools J, DeAngelo DJ, Gotlib J *et al.* A tyrosine kinase created by fusion of the PDGFRA and FIP1L1 genes as a therapeutic target of imatinib in idiopathic hypereosinophilic syndrome. *N Engl J Med* 2003; **348**:1201–14.
- 82 Pardanani A, Ketterling RP, Brockman SR *et al.* CHIC2 deletion, a surrogate for FIP1L1–PDGFRA fusion, occurs in systemic mastocytosis associated with eosinophilia and predicts response to imatinib mesylate therapy. *Blood* 2003; **102**:3093–6.
- 83 Robyn J, Lemery S, McCoy JP *et al.* Multilineage involvement of the fusion gene in patients with FIP1L1/PDGFRA-positive hypereosinophilic syndrome. *Br J Haematol* 2006; **132**:286–92.
- 84 Gotlib J, Cross NCR, Gilliland DG. Eosinophilic disorders: molecular pathogenesis, new classification, and modern therapy. *Best Pract Res Clin Haematol* 2006; **19**:535–69.
- 85 Wilson TM, Maric I, Simakova O *et al.* Clonal analysis of NRAS activating mutations in KIT-D816V systemic mastocytosis. *Haematologica* 2011; **96**:459–63.
- 86 Bandi SR, Brandts C, Rensinghoff M *et al.* E3 ligase-defective Cbl mutants lead to a generalized mastocytosis and a myeloproliferative disease. *Blood* 2009; **114**:4197–208.
- 87 van Rijn RM, van Marle A, Chazot PL *et al.* Cloning and characterization of dominant negative splice variants of the human histamine H4 receptor. *Biochem J* 2008; **414**:121–31.
- 88 Tefferi A, Levine RL, Lim KH *et al.* Frequent TET2 mutations in systemic mastocytosis: clinical, KITD816V and FIP1L1–PDGFRA correlates. *Leukemia* 2009; **23**:900–4.
- 89 Schroer BC, Han YC, Hsieh FH. Identification of TET2 mutations in subjects with systemic mastocytosis and hypereosinophilic syndrome[abstract]. *J Allergy Clin Immunol* 2010; **125**:AB232.
- 90 Nedoszytko B, Nedoszytko M, Lange M *et al.* Associations of TNF- α gene polymorphisms and mastocytosis. A study of the European Competence Network in Mastocytosis (ECNM)[abstract]. *Allergy* 2008; **63**(Suppl. 88):154.
- 91 Nedoszytko M, Nedoszytko B, Lange M *et al.* Vascular Endothelial Growth Factor (VEGF) promoter gene polymorphism in mastocytosis. A study of the European Competence Network on Mastocytosis (ECNM)[abstract]. *Allergy* 2008; **63**(Suppl. 88):322.
- 92 Alfter K, von Kügelgen I, Haenisch B *et al.* New aspects of liver abnormalities as part of the systemic mast cell activation syndrome. *Liver Int* 2009; **29**:181–6.